

**ROWLEY  
BIOCHEMICAL  
INC.**

**DANVERS INDUSTRIAL PARK  
10 ELECTRONICS AVENUE,  
DANVERS MA 01923  
TEL: 978-739-4883 FAX: 978-739-5640  
www.rowleybio.com**

---

## **G-473 TOLUIDINE BLUE FOR MAST CELLS**

**FIXATION:** 10% Buffered Neutral Formalin (F-113)

**SECTION:** Paraffin, @ 4 to 5um

### **STAINING PROCEDURE:**

1. Deparaffinize and hydrate to distilled water.
2. Stain sections in Toluidine Blue, 1% Aqueous (G-473-1) for 10 minutes.
3. Rinse in distilled water.
4. Quickly dehydrate through 95% and absolute alcohols.
5. Clear in Xylene and mount with synthetic resin.

### **NOTES:**

1. Alcohol cannot be used for dehydration in most methachromatic staining procedures, but metachromasia of mast cell granules are stable and will not be lost after alcoholic dehydration.
2. A rapid screening method for mast cells is to simply apply a light Methylene Blue stain as if you were counterstaining an acid-fast stain. The mast cells are very well demonstrated.

### **REFERENCES:**

Carson F.L., Histotechnology, A Self-instructional Text. ASCP Chicago.1990, p.163-164.