ROWLEY
BIOCHEMICAL
INC.

DANVERS INDUSTRIAL PARK
10 ELECTRONICS AVENUE,
DANVERS MA 01923
TEL: 978-739-4883 FAX: 978-739-5640
www.rowleybio.com

L-755, HEIDENHAIN'S HEMATOXYLIN

Fixation: **Zenker (F1-55), Bouin's(F-40), or Helly (F-153)**

Sections: Embedded in Parrafin

Staining:

1. Deparaffinize and hydrate to distilled water.
2. Mordant in **Ferric Ammonium Sulfate, 4%** (L-755-2) for 30 minutes to 3 hours.
3. Stain 1-3 hours in **Heidenhain's Hematoxylin** (L-755-1). Rinse in tap water.
4. Differentiate in **Ferric Ammonium Sulfate, 4%** (L-755-2). Control differentiation by microscopic examination. Wash in running water 10 minutes.
5. Counterstain, if desired.
6. Dehydrate, clear and mount on **Canada Balsam** (M-6).

Stain Results:

(Nuclei)	Black

References:

Clark, G.: Staining Procedures, Williams and Wilkins Co., Baltimore, 3rd Ed., c. 1973, p.38

Conn, H.J.: Biological Stains, 8th Ed., Williams and Wilkins Co., Baltimore, 1969, p. 437.